

Table 8. No till edible dry bean establishment and yield when seeded into a winter rye companion crop.

TRT	Edible Bean	Rye Termination		Population	Weed Biomass at Harvest			Yield	
		Date	lb/acre	plants/acre	Grass lb/acre	Broadleaf lb/acre	Total lb/acre	(bu/acre)	
May 1 Kill and till ¹	Zorro	1-May	645	115434	37	17	54	50	a
May 1 Kill and till ¹	Majesty	1-May	645	84398	55	130	185	33	cd
May 1 Kill and till ¹	Tiger's Eye	1-May	645	92565	116	29	144	22	efg
Boot stage Chop and till ²	Zorro	2-Jun	2477	116795	53	51	103	42	b
Boot stage Chop and till ²	Majesty	2-Jun	2477	93110	29	26	55	26	def
Boot stage Chop and till ²	Tiger's Eye	2-Jun	2477	95288	22	25	47	19	fg
Anthesis Chop and till ³	Zorro	10-Jun	3287	108628	44	13	57	35	bc
Anthesis Chop and till ³	Majesty	10-Jun	3287	96377	35	5	39	29	cde
Anthesis Chop and till ³	Tiger's Eye	10-Jun	3287	94743	74	17	91	16	g
Anthesis roll ⁴	Zorro	10-Jun	3287	105089	148	355	503	30	cd
Anthesis roll ⁴	Majesty	10-Jun	3287	81947	166	259	425	22	efg
Anthesis roll ⁴	Tiger's Eye	10-Jun	3287	86303	61	182	244	15	g

¹Power Harrow on 5/1/14. Disk and Power Harrow on 6/17/14. Row cultivation on 7/15/14.

²Disk and Power Harrow on 6/2/14. Disk and Power Harrow on 6/17/14. Row cultivation on 7/15/14.

³Chop and Power Harrow on 6/10/14. Disk and Power Harrow on 6/17/14. Row cultivation on 7/15/14.

⁴Roll on 6/10/14 (3X). Roll on 6/17/14 (3X). Row cultivation on 7/15/14.

Within a column for a given set of treatments, means followed by the same letter are not significantly different at (α 0.05).