

Table 3. Comparison of treatment methods for implementation into existing cranberry beds.

<b>Treatment</b>	<b>Cost/ foot</b>	<b>Advantages</b>	<b>Disadvantages</b>	
Under drain	.80	No effect on harvest	Plastic pipe placed in ground.	Highest cost
Deep homogenization	.25	No loss of area. Inexpensive	May work under limited conditions	
Ditches	.15	Inexpensive	Annual maintenance	Disrupts harvest

**References**

- Caruso, F. L. & Wilcox, W. F. (1990). *Phytophthora cinnamomi* as a cause of root rot and dieback of cranberry in Massachusetts. *Plant Disease* **74**, 664-667.
- Erwin, D. C. & Ribero, O. K. (1996). *Phytophthora diseases worldwide*. St. Paul: APS Press.
- Oudemans, P. V. (1995). *Phytophthora* root rot of cranberries: identification of inoculum sources by lupin baiting. *Phytopathology* **85**, abstract.
- Oudemans, P. V. (1998). Detection and monitoring of *Phytophthora cinnamomi* in cranberry irrigation water sources via lupin baiting. *Plant Disease* **82** (in press).
- Zentmyer, G. A. (1980). *Phytophthora cinnamomi* and the diseases it causes. St. Paul: The American Phytopathological Society.